Recreational fishing profile 2020/21

South Coast Bioregion

Here's some data about the South Coast Bioregion that recreational fishers helped generate through our Statewide Survey of Recreational Fishing.

The South Coast Bioregion of Western Australia spans from Black Point (east of Augusta) to the WA/SA border, including the regional centres of Albany and Esperance. The most frequently caught species in this bioregion were King George whiting, Australian herring and school whiting. The most frequently caught invertebrate species was squid.



Catch the facts on recreational fishing in the South Coast Bioregion



13%

of Recreational Fishing from Boat Licence (RFBL) holders reported fishing in the bioregion



6%

of days fished in WA were in this bioregion



27,325

fishing events



94,839

hours fished



25,886

boat days





8%

of statewide fishing effort (hours)

92

species reported





To learn more about statewide recreational fishing, visit our webpage: fish.wa.gov.au/statewide-recfishing

Important disclaimer

The Chief Executive Officer of the Department of Primary Industries and Regional Development and the State of Western Australia accept no liability whatsoever by reason of negligence or otherwise arising from the use or release of this information or any part of it.

Copyright © State of Western Australia (Department of Primary Industries and Regional Development), 2023

Department of Primary Industries and Regional Development

+61 1300 374 731 | research.survey@dpird.wa.gov.au | dpird.wa.gov.au

ABN: 18 951 343 745